

Allergy Care Plan Request Form

Child's name: _____

Child's date of birth: _____

The child listed above attends our child care or early learning program. We have been informed that they have been diagnosed with an allergy.

Child Care Program Director: _____

Child Care Program: _____

Mailing Address: _____

Phone Number: _____

Fax Number: _____

Healthcare Provider: As a licensed child care program, we are required to meet state licensing standards (WAC 110-300-0215 and 110-300-0300). Please complete the following Allergy and Anaphylaxis Emergency Plan and, if necessary, a Medication Authorization Form. We need to know what the child is allergic to, the steps to take in response to a suspected allergic reaction, and any relevant food substitutions.

By signing below, I give permission to my child's healthcare provider to release the information requested above to my child care program.

Parent or Guardian Name (Printed): _____

Parent or Guardian Signature: _____

Date: _____

Parent or Guardian Phone Number: _____

Allergy and Anaphylaxis Emergency Plan

American Academy of Pediatrics


DEDICATED TO THE HEALTH OF ALL CHILDREN®



Child's name: _____ Date of plan: _____

Date of birth: ____/____/____ Age ____ Weight: _____kg

Child has allergy to _____



- Child has asthma. Yes No (If yes, higher chance severe reaction)
Child has had anaphylaxis. Yes No
Child may carry medicine. Yes No
Child may give him/herself medicine. Yes No (If child refuses/is unable to self-treat, an adult must give medicine)

IMPORTANT REMINDER

Anaphylaxis is a potentially life-threatening, severe allergic reaction. If in doubt, give epinephrine.

For Severe Allergy and Anaphylaxis What to look for



If child has ANY of these severe symptoms after eating the food or having a sting, **give epinephrine.**

- Shortness of breath, wheezing, or coughing
- Skin color is pale or has a bluish color
- Weak pulse
- Fainting or dizziness
- Tight or hoarse throat
- Trouble breathing or swallowing
- Swelling of lips or tongue that bother breathing
- Vomiting or diarrhea (if severe or combined with other symptoms)
- Many hives or redness over body
- Feeling of "doom," confusion, altered consciousness, or agitation

SPECIAL SITUATION: If this box is checked, child has an extremely severe allergy to an insect sting or the following food(s): _____. Even if child has MILD symptoms after a sting or eating these foods, **give epinephrine.**

Give epinephrine! What to do

1. Inject epinephrine right away! Note time when epinephrine was given.
2. Call 911.
 - Ask for ambulance with epinephrine.
 - Tell rescue squad when epinephrine was given.
3. Stay with child and:
 - Call parents and child's doctor.
 - Give a second dose of epinephrine, if symptoms get worse, continue, or do not get better in 5 minutes.
 - Keep child lying on back. If the child vomits or has trouble breathing, keep child lying on his or her side.
4. Give other medicine, if prescribed. Do not use other medicine in place of epinephrine.
 - Antihistamine
 - Inhaler/bronchodilator

For Mild Allergic Reaction What to look for



If child has had any mild symptoms, **monitor child.**

Symptoms may include:

- Itchy nose, sneezing, itchy mouth
- A few hives
- Mild stomach nausea or discomfort

Monitor child What to do

Stay with child and:

- Watch child closely.
- Give antihistamine (if prescribed).
- Call parents and child's doctor.
- If more than 1 symptom or symptoms of severe allergy/anaphylaxis develop, use epinephrine. (See "For Severe Allergy and Anaphylaxis.")

Medicines/Doses

Epinephrine, intramuscular (list type): _____ Dose: 0.10 mg (7.5 kg to less than 13 kg)*
 0.15 mg (13 kg to less than 25 kg)
 0.30 mg (25 kg or more)

Antihistamine, by mouth (type and dose): _____ (*Use 0.15 mg, if 0.10 mg is not available)

Other (for example, inhaler/bronchodilator if child has asthma): _____

Parent/Guardian Authorization Signature

Date

Physician/HCP Authorization Signature

Date

Allergy and Anaphylaxis Emergency Plan

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®



Child's name: _____ Date of plan: _____

Additional Instructions:

Contacts

Call 911 / Rescue squad: _____

Doctor: _____ Phone: _____

Parent/Guardian: _____ Phone: _____

Parent/Guardian: _____ Phone: _____

Other Emergency Contacts

Name/Relationship: _____ Phone: _____


Name/Relationship: _____ Phone: _____

There are many different types of Epinephrine auto-injectors. **Always follow the instructions on the medication label, as well as any child-specific instructions from parents or guardians.**

Below are common types of epinephrine auto-injectors and how to use them. **These instructions* DO NOT replace staff training by the parent or guardian.**


HOW TO USE AUVI-Q® (EPINEPHRINE INJECTION, USP), KALEO

1. Remove Auvi-Q from the outer case. Pull off red safety guard.
2. Place black end of Auvi-Q against the middle of the outer thigh.
3. Press firmly until you hear a click and hiss sound, and hold in place for 2 seconds.
4. Call 911 and get emergency medical help right away.




HOW TO USE EPIPEN®, EPIPEN JR® (EPINEPHRINE) AUTO-INJECTOR AND EPINEPHRINE INJECTION (AUTHORIZED GENERIC OF EPIPEN®), USP AUTO-INJECTOR, MYLAN AUTO-INJECTOR, MYLAN

1. Remove the EpiPen® or EpiPen Jr® Auto-Injector from the clear carrier tube.
2. Grasp the auto-injector in your fist with the orange tip (needle end) pointing downward. With your other hand, remove the blue safety release by pulling straight up.
3. Swing and push the auto-injector firmly into the middle of the outer thigh until it 'clicks'. Hold firmly in place for 3 seconds (count slowly 1, 2, 3).
4. Remove and massage the injection area for 10 seconds. Call 911 and get emergency medical help right away.



HOW TO USE IMPAX EPINEPHRINE INJECTION (AUTHORIZED GENERIC OF ADRENALCLICK®), USP AUTO-INJECTOR, AMNEAL PHARMACEUTICALS

1. Remove epinephrine auto-injector from its protective carrying case.
2. Pull off both blue end caps: you will now see a red tip. Grasp the auto-injector in your fist with the red tip pointing downward.
3. Put the red tip against the middle of the outer thigh at a 90-degree angle, perpendicular to the thigh. Press down hard and hold firmly against the thigh for approximately 10 seconds.
4. Remove and massage the area for 10 seconds. Call 911 and get emergency medical help right away.



HOW TO USE TEVA'S GENERIC EPIPEN® (EPINEPHRINE INJECTION, USP) AUTO-INJECTOR, TEVA PHARMACEUTICAL INDUSTRIES

1. Quickly twist the yellow or green cap off of the auto-injector in the direction of the "twist arrow" to remove it.
2. Grasp the auto-injector in your fist with the orange tip (needle end) pointing downward. With your other hand, pull off the blue safety release.
3. Place the orange tip against the middle of the outer thigh at a right angle to the thigh.
4. Swing and push the auto-injector firmly into the middle of the outer thigh until it 'clicks'. Hold firmly in place for 3 seconds (count slowly 1, 2, 3).
5. Remove and massage the injection area for 10 seconds. Call 911 and get emergency medical help right away.



HOW TO USE SYMJEPI™ (EPINEPHRINE INJECTION, USP)

1. When ready to inject, pull off cap to expose needle. Do not put finger on top of the device.
2. Hold SYMJEPI by finger grips only and slowly insert the needle into the thigh. SYMJEPI can be injected through clothing if necessary.
3. After needle is in thigh, push the plunger all the way down until it clicks and hold for 2 seconds.
4. Remove the syringe and massage the injection area for 10 seconds. Call 911 and get emergency medical help right away.
5. Once the injection has been administered, using one hand with fingers behind the needle slide safety guard over needle.



ADMINISTRATION AND SAFETY INFORMATION FOR ALL AUTO-INJECTORS:

1. Do not put your thumb, fingers or hand over the tip of the auto-injector or inject into any body part other than mid-outer thigh. In case of accidental injection, go immediately to the nearest emergency room.
2. If administering to a young child, hold their leg firmly in place before and during injection to prevent injuries.
3. Epinephrine can be injected through clothing if needed.
4. Call 911 immediately after injection.